

iMEC™ with TAPE ORAL BROAD SPECTRUM WORMER AND BOTICIDE FOR HORSES

ACTIVE CONSTITUENTS: 15.5mg/g IVERMECTIN, 77.5mg/g PRAZIQUANTEL

Easy worming with less paste

For the treatment and control of intestinal round worms (including larval stages of *strongylus vulgaris* in intestinal arteries and benzimidazole resistant strongyles), lungworms, tapeworms, bots, and skin lesions caused by *Habronema* and *Draschia* spp.(summer sores) and *onchocerca* spp. microfilariae (cutaneous onchocerciasis).

- Low volume
- Easy to administer paste formulation
- Safe to use in all horses
- Cost effective



	iMEC™ with TAPE	EQUIMEC® Tape
Active Constituents	15.5mg/g IVERMECTIN 77.5mg/g PRAZIQUANTEL	15.5mg/g IVERMECTIN 77.5mg/g PRAZIQUANTEL
Species	Horses	Horses
Route of Administration	Oral	Oral
Syringe size	7.74g	7.74g
Dose rate	1 tube per 600kg	1 tube per 600kg

iMEC™ is a trademark of Abbey Laboratories Pty Ltd

EQUIMEC® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH

iMEC™ with TAPE ORAL BROAD SPECTRUM WORMER AND BOTICIDE FOR HORSES

ACTIVE CONSTITUENTS: 15.5mg/g IVERMECTIN, 77.5mg/g PRAZIQUANTEL



For the treatment and control of intestinal round worms (including larval stages of *strongylus vulgaris* in intestinal arteries and benzimidazole resistant strongyles), lungworms, tapeworms, bots, and skin lesions caused by *Habronema* and *Draschia* spp.(summer sores) and *onchocerca* spp. *microfilariae* (cutaneous onchocerciasis).

INDICATIONS

At the recommended dose level of 200µg ivermectin and 1mg/kg praziquantel liveweight, this product controls the following parasites of horses:

TAPEWORMS

Anoplocephala perfoliata

LARGE STRONGYLES

Strongylus vulgaris (adults, arterial larval stages),
Strongylus edentatus (adults and tissue stages),
Strongylus equinus (adults),
Triodontophorus spp. (adults) including *T.brevicauda*, *T.serratus* and *T.tenuicollis* and *Craterostomum acuticaudatum* (adults)

SMALL STRONGYLES

Including benzimidazole resistant strains of adult and immature:

Cyathostomum spp, *Cyath.catinatum* and *Cyath.pateratum*; *Cylicocycclus* spp including *Cylicocyc.ashworthi*, *Cylicocyc.elongatus*, *Cylicocyc.insigne*, *Cylicocyc.leptostomum*, *Cylicocyc.nassatus*, *Cylicocyc.radiatus*; *Cylicostephanu* spp. including *Cylicost asymentricus*, *Cylicost. bidentatus*, *Cylicost.calicatus*, *Cylicost. goldi*, *Cylicost.longibursatus*, *Cylicost. minutus*, *Cylicodontophorus* spp. including *Cylicodontophorus bicoronatus*; *Coronocycclus* spp., including *Co. coronatus*, *Co.labiatus*, *Co.labratus*; *Gyalcephalus* spp. including *G.capitatus*, *Parapoteriostomum* spp. including *Parap.euproctus* and *Parap. mettamis*, *Petrovinema* spp. including *Petr.poculatum* and *Poteriostomum* spp. including *Pot.imparidentatum* and *Pot.ratzii*

PINWORMS

Oxyuris equi (adult and immature)

ASCARIDS

Parascaris equorum (adult and immature)

HAIRWORMS

Trichostrongylus axei (adult)

LARGE MOUTHED STOMACH WORMS

Habronema muscae (adult)

NECK THREADWORM

Onchocerca spp (*microfilariae*).

BOTS

Gasterophilus intestinalis and *Gasterophilus nasalis* (oral, gastric and intestinal stages)

LUNGWORMS

Dictyocaulus amfieldi (adult and immature)

INTESTINAL THREADWORM

Strongyloides westeri (adult)

Resistance may develop to any chemical.

DIRECTIONS FOR USE

SIDE EFFECTS

Some horses have experienced reactions with swelling and itching shortly after treatment. In most cases the horses were diagnosed as carrying heavy infections of *Onchocerca microfilariae* and it is assumed the reactions were the result of *microfilariae* dying in large numbers. Although the signs may resolve in a few days, symptomatic treatment may be advisable. Consult your veterinarian if these signs persist.

DOSAGE AND ADMINISTRATION

iMEC™ with TAPE BROAD SPECTRUM ANTIPARASITIC PASTE FOR HORSES is given orally at the recommended dose level of 200µg ivermectin and 1mg/kg praziquantel liveweight. **The contents of this syringe will treat one horse of 600kg liveweight. Each weight marking on the syringe plunger will deliver 1.29g of paste which is sufficient to treat 100kg live weight.** Remove the plastic cap from the tip of the nozzle. Make sure the horse's mouth does not contain feed. Insert the syringe into the side of horse's mouth in the gap between the front and back teeth. Advance the plunger as far as it will go depositing the paste on the rear of the tongue. Immediately raise the horse's head for few seconds after dosing to ensure the full dose has been swallowed.

GENERAL DIRECTIONS

PARASITE CONTROL PROGRAMS

All horses should be included in a regular parasite control program with particular attention paid to mares foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. iMEC™ with TAPE is highly effective against gastrointestinal cutaneous and pulmonary nematodes, tapeworms and bots of horses. Regular treatment will reduce the chances of verminous arteritis and colic caused by *S.vulgaris*.

With its broad spectrum activity, iMEC™ with TAPE is well suited to be the major product in a parasite control program. Consult your veterinarian for control program to meet your specific needs.

PRODUCT ADVANTAGES

BROAD SPECTRUM

iMEC™ with TAPE kills all the important internal parasites of horses, including the arterial stages of *S.vulgaris* and bots with a single, low volume, easy to administer dose. Because reinfection is common, treatment must be repeated routinely for effective control.

SAFETY TO HORSES

Ivermectin and Praziquantel have wide safety margin at recommended dose level. They may be used in horses of all ages. Mares may be treated at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

Resolution of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with iMEC™ with TAPE. Reinfection and measures for it's prevention should also be considered. Consult your veterinarian if the condition does not improve.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

MEAT WITHHOLDING PERIOD

DO NOT USE less than 28 days before slaughter for human consumption.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

ENVIRONMENTAL PROTECTION

Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Dispose of container by wrapping with paper and putting in the garbage.

STORAGE

Store below 30°C (Room Temperature). Protect from light.

APVMA Approval Number: 92178/137750

iMEC™ is a trademark of Abbey Laboratories Pty Ltd